



The biodiversity of aerobic endospore-forming bacterial species occurring in Yanyanku and Ikpiru, fermented seeds of *Hibiscus sabdariffa*

Agbobatinkpo, Pélagie Bidossessi; Thorsen, Line; Nielsen, Dennis Sandris; Azokpota, Paulin; Akissoe, Noël; Hounhouigan, Joseph; Jakobsen, Mogens

Published in:
Global issues in food microbiology

Publication date:
2012

Document version
Publisher's PDF, also known as Version of record

Citation for published version (APA):
Agbobatinkpo, P. B., Thorsen, L., Nielsen, D. S., Azokpota, P., Akissoe, N., Hounhouigan, J., & Jakobsen, M. (2012). The biodiversity of aerobic endospore-forming bacterial species occurring in Yanyanku and Ikpiru, fermented seeds of *Hibiscus sabdariffa*. In *Global issues in food microbiology: abstract book* (pp. 659)





23rd INTERNATIONAL ICFMH SYMPOSIUM
FoodMicro 2012



Global Issues In Food Microbiology

Abstract Book



FoodMicro

3-7 SEPTEMBER 2012 ISTANBUL

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

Food products

RISTIMUÑO², Pier Sandro
sa AZNAR¹
Burjasot, SPAIN

di Microbiologia-Centro Ricerche

beverage prepared from maize in the
acteria (LAB) species present during
analyzed by molecular methods.
mentation process using maize flour
rst step, manually shaped buns were
ie sugar, water and fermented (18-20
d with fresh maize flour and water,
is were separated before the final
ial counts were in the range of 10⁴-
mong assayed media. A total of 146
ing steps were identified by internal
orphic DNA (RAPD) profiles as
1 (20%), *Lactobacillus* (23%) and
species-specific PCR revealed *E.*
lactis and *W. viridescens* as the
uction process showed the presence
ing exhibited the highest diversity
roides and *E. mundtii*. From pre-
b. brevis and *Leuc. mesenteroides*
d as the dominant species in the
ely *Leuc. lactis* while the jelly-like
ng concentration, both species but
les revealed intraspecies diversity,
motolerant *Enterococcus faecium*
Leuc. mesenteroides (13%) that
indigenous foods may provide
which can be exploited as new
nal quality of traditional Andean

P-471

The biodiversity of aerobic endospore-forming bacterial species occurring in Yanyanku and Ikpiru, fermented seeds of *Hibiscus sabdariffa*

Pélagie Bidossessi AGBOBATINKPO¹, Line THORSEN², Dennis Sandris
NIELSEN², Paulin AZOKPOTA¹, Noël AKISSOE¹, Joseph
HOUNHOUIGAN¹, Mogens JAKOBSEN²

1: University of Abomey-calavi (Benin), Faculty of Agricultural Sciences, Nutrition and Food Sciences, BENIN

2: University of Copenhagen, Food Science, Faculty of Life Sciences, DENMARK

Corresponding author: abpelagic@yahoo.fr

Yanyanku and Ikpiru are two fermented products from Malcavene bean (*Hibiscus sabdariffa*) used as functional additives for African locust bean seeds (*Parkia biglobosa*) fermentation in Benin. In the present study, a total of 343 Aerobic Endospore-Forming Bacteria (AEFB) isolated from Yanyanku and Ikpiru produced in northern and southern Benin, at different production sites were identified using phenotypic and molecular biology based methods, including 16S rRNA, *gyrA* and *gyrB* genes sequencing. Detection of genes encoding cytotoxin K (cytK1, cytK2), haemolysin BL (hblA, hblC, hblD) and a genetic determinant for the emetic toxin cereulide EM1 were also performed. Regardless of geographical location and production site, it was the same five-six predominant species that were involved in the fermentation of *H. sabdariffa* for the production of Yanyanku and Ikpiru. The predominant species were of the genus *Bacillus* and included *B. subtilis* (19-41% of isolates), *B. cereus* (6-41%), *B. amyloliquefaciens* (9-20%), *B. safensis* (6-23%), *B. licheniformis* (4-25%), and *B. altitudinis* (0-19%). Other species occurred sporadically and included *B. flexus*, *B. circulans* (<1% each), as well as species of the genera (5.5% of isolates) *Lysinibacillus*, *Brevibacillus*, *Aneurinibacillus* and *Paenibacillus*. Sequencing of the *gyrA* gene showed to be an efficient marker for differentiation of *B. pumilus* and *B. safensis*. All the *B. cereus* isolates lacked the gene encoding the cytotoxin K-1 but 91% of them harboured the gene encoding the cytotoxin K-2 and 6% the emetic specific gene fragment EM1. The genes encoding haemolysin BL hblA, hblC, hblD were present in 15%, 34% and 35% of *B. cereus* isolates, respectively. None of *B. cereus* from functional additives Yanyanku and Ikpiru harboured all the toxin genes investigated. This study is the first to identify the AEFB of the functional additives Yanyanku and Ikpiru to species level and perform a safety evaluation based on toxin gene detections. The results can be used to initiate selection of suitable startercultures for the controlled fermentation of Yanyanku and Ikpiru in order to provide a stable and safe product.